# Optimization for Machine Learning 机器学习中的优化方法

#### 陈程

华东师范大学 软件工程学院

chchen@sei.ecnu.edu.cn

#### Outline

Review

2 Projected Gradient Descent

3 Frank-Wolfe Algorithm

# Review of Smooth and Strongly Convex

A differentiable function f is L-smooth if

$$f(\mathbf{y}) \le f(\mathbf{x}) + \langle \nabla f(\mathbf{x}), \mathbf{y} - \mathbf{x} \rangle + \frac{L}{2} \|\mathbf{x} - \mathbf{y}\|_2^2, \ \forall \ \mathbf{x}, \mathbf{y} \in \mathbb{R}^d$$

A differentiable function f is  $\mu$ -strongly convex if

$$f(\mathbf{y}) \ge f(\mathbf{x}) + \langle \nabla f(\mathbf{x}), \mathbf{y} - \mathbf{x} \rangle + \frac{\mu}{2} \|\mathbf{x} - \mathbf{y}\|_2^2, \ \forall \ \mathbf{x}, \mathbf{y} \in \mathbb{R}^d$$

 $\kappa \triangleq \frac{L}{\mu}$  is the condition number.

 Lecture 05
 OptML
 November 1st, 2023
 2 / 26

#### Review of Gradient Descent

Consider an unconstrained convex optimization problem

$$\min_{\mathbf{x}\in\mathbb{R}^d} f(\mathbf{x}).$$

The **gradient descent** method starts with an initial point  $\mathbf{x}_0$ , and iteratively computes

$$\mathbf{x}_{t+1} = \mathbf{x}_t - \eta_t \nabla f(\mathbf{x}_t).$$

#### Review of Gradient Descent

	stepsize	convergence rate	iteration complexity
strongly convex & smooth	$\eta_t = rac{1}{L}  ext{ or } \ \eta_t = rac{2}{\mu + L}$	$O\left(\left(1-\frac{1}{\kappa}\right)^t\right)$	$O(\kappa \log \frac{1}{\varepsilon})$
locally strongly convex & smooth	$\eta_t = \frac{1}{L}$	$O\left(\left(1-\frac{1}{\kappa}\right)^t\right)$	$O(\kappa \log \frac{1}{\varepsilon})$
PL condition & smooth	$\eta_t = \frac{1}{L}$	$O\left(\left(1-\frac{1}{\kappa}\right)^t\right)$	$O(\kappa \log \frac{1}{\varepsilon})$
convex & smooth	$\eta_t = \frac{1}{L}$	$O\left(\frac{1}{t}\right)$	$O\left(\frac{1}{\varepsilon}\right)$
nonconvex & smooth	$\eta_t = \frac{1}{L}$	$O\left(\frac{1}{\sqrt{t}}\right)$	$O\left(\frac{1}{\varepsilon^2}\right)$

Table: Convergence Property of GD

 Lecture 05
 OptML
 November 1st, 2023
 4 / 26

#### Outline

Review

2 Projected Gradient Descent


Frank-Wolfe Algorithm

#### **Constrained Convex Optimization**

Suppose f is a convex function and  $C \in \mathbb{R}^d$  is a closed and convex set. The constrained convex optimization problem is:

$$\min_{\mathbf{x}} f(\mathbf{x})$$
  
s.t.  $\mathbf{x} \in C$ 

# **Optimality Condition**




Suppose f is convex and differentiable. Then

$$\mathbf{x}^* \in \operatorname*{arg\,min}_{\mathbf{x} \in \mathcal{C}} f(\mathbf{x}) \iff \langle -\nabla f(\mathbf{x}^*), \mathbf{z} - \mathbf{x}^* \rangle \leq 0, \ \forall \ \mathbf{z} \in \mathcal{C}$$

 Lecture 05
 OptML
 November 1st, 2023
 6 / 26

# Projected Gradient Descent (投影梯度下降法)




Idea: project onto  $\mathcal C$  after every gradient descent step:

$$\mathbf{x}_{t+1} = \mathcal{P}_{\mathcal{C}}(\mathbf{x}_t - \eta_t \nabla f(\mathbf{x}_t)).$$

where  $\mathcal{P}_{\mathcal{C}}(\mathbf{x}) \triangleq \arg\min_{\mathbf{z} \in \mathcal{C}} \|\mathbf{z} - \mathbf{x}\|_{2}^{2}$  is Euclidean projection onto  $\mathcal{C}$ .

 Lecture 05
 OptML
 November 1st, 2023
 7 / 26


# Properties of Projection



Let  $\mathcal{C} \in \mathbb{R}^d$  be closed and convex,  $\mathbf{z} \in \mathcal{C}$ ,  $\mathbf{x} \in \mathbb{R}^d$ . Then

 Lecture 05
 OptML
 November 1st, 2023
 8 / 26

# Properties of Projection: Nonexpansivness



Let  $\mathcal{C} \in \mathbb{R}^d$  be closed and convex. For any  $\mathbf{x}, \mathbf{z} \in \mathbb{R}^d$ , we have

$$\|\mathcal{P}_{\mathcal{C}}(\mathbf{x}) - \mathcal{P}_{\mathcal{C}}(\mathbf{z})\|_2 \le \|\mathbf{x} - \mathbf{z}\|_2$$
.

 Lecture 05
 OptML
 November 1st, 2023
 9 / 26

#### **Think**

Suppose f is a convex function and C is a closed convex set. Let


$$\hat{\mathbf{x}} = \underset{\mathbf{x} \in \mathbb{R}^d}{\min} \ f(\mathbf{x}) \ \text{and} \ \mathbf{x}^* = \underset{\mathbf{x} \in \mathcal{C}}{\arg\min} \ f(\mathbf{x})$$

Is it true that

$$\textbf{x}^* = \mathcal{P}_{\mathcal{C}}(\hat{\textbf{x}})?$$

10 / 26

# **Examples**




Projecting onto an affine subspace:

$$\begin{aligned} \mathbf{y} &= \operatorname*{arg\,min}_{\mathbf{z}} \left\| \mathbf{A} \mathbf{z} - \mathbf{x} \right\|_2 = (\mathbf{A}^{\top} \mathbf{A})^{-1} \mathbf{A}^{\top} \mathbf{x} \\ \\ \mathcal{P}_{\mathbf{A}}(\mathbf{x}) &= \mathbf{A} \mathbf{y} = \mathbf{A} (\mathbf{A}^{\top} \mathbf{A})^{-1} \mathbf{A}^{\top} \mathbf{x} \end{aligned}$$

 Lecture 05
 OptML
 November 1st, 2023
 11/26

# **Examples**




Projecting onto a unit Euclidean ball ( $\ell_2$  ball):

$$\mathcal{P}_{\mathcal{C}}(\mathbf{x}) = \mathop{\arg\min}_{\|\mathbf{z}\|_2 \leq 1} \|\mathbf{x} - \mathbf{z}\|_2 = \frac{\mathbf{x}}{\max\{1, \|\mathbf{x}\|_2\}}$$

Lecture 05 OptML November 1st, 2023 12 / 26

#### Examples



Projecting onto a unit  $\ell_1$  ball:

$$\mathbf{y} = \mathcal{P}_{\mathcal{C}}(\mathbf{x}) = \mathop{\arg\min}_{\|\mathbf{z}\|_1 \leq 1} \|\mathbf{x} - \mathbf{z}\|_2$$

If  $\|\mathbf{x}\|_1 \leq 1$  then  $\mathcal{P}_{\mathcal{C}}(\mathbf{x}) = \mathbf{x}$ . Otherwise,

$$y_i = sign(x_i)(|x_i| - \lambda)_+$$

where  $(\cdot)_+ = \max\{\cdot, 0\}$  and  $\lambda$  is the root of  $\sum_{i=1}^n (|x_i| - \lambda)_+ = 1$ .

Lecture 05 OptML Novem

13 / 26

# Smooth and Strongly Convex Constrained Problems

$$\min_{\mathbf{x}} f(\mathbf{x})$$
  
s.t.  $\mathbf{x} \in \mathcal{C}$ 

- f: L-smooth and  $\mu$ -strongly convex
- ullet  $\mathcal{C} \in \mathbb{R}^d$ : closed and convex

# Smooth and Strongly Convex Constrained Problems

Let f be L-smooth and  $\mu$ -strongly convex. If  $\eta_t = \eta = \frac{2}{\mu + L}$ , then PGD obeys

$$\left\|\mathbf{x}_{t}-\mathbf{x}^{*}\right\|_{2} \leq \left(rac{\kappa-1}{\kappa+1}
ight)^{t}\left\|\mathbf{x}_{0}-\mathbf{x}^{*}\right\|_{2}.$$

the same convergence rate as for the unconstrained case

Lecture 05 OptML November 1st, 2023 15 / 26

# Contraction Mapping (压缩映射)

**Contraction mapping in Euclidean space**: If a function  $f: \mathcal{X} \to \mathcal{X}$  satisfies

$$||f(x) - f(y)||_2 \le \gamma ||x - y||_2, \ \forall \ x, y \in \mathcal{X}$$

for some  $\gamma \in (0,1)$ , then we call f is a contraction mapping.

The contraction mapping f has a unique fixed point  $\hat{x}$ , i.e.,  $f(\hat{x}) = \hat{x}$ .

 Lecture 05
 OptML
 November 1st, 2023
 16 / 26

#### Smooth and Convex Constrained Problems

$$\min_{\mathbf{x}} f(\mathbf{x})$$
  
s.t.  $\mathbf{x} \in C$ 

- f: convex and L-smooth
- ullet  $\mathcal{C} \in \mathbb{R}^d$ : closed and convex

#### Smooth and Convex Constrained Problems

Let f be convex and L-smooth. If  $\eta_t = \eta = \frac{1}{L}$ , then PGD obeys

$$f(\mathbf{x}_t) - f(\mathbf{x}^*) \le \frac{2L\|\mathbf{x}_0 - \mathbf{x}^*\|^2}{t}$$

the same convergence rate as for the unconstrained case

 Lecture 05
 OptML
 November 1st, 2023
 18 / 26

# Convergence Analysis

Recall the main steps when handling the unconstrained case:

• Step 1: show improvement

$$f(\mathbf{x}_{t+1}) \leq f(\mathbf{x}_t) - \frac{1}{2I} \|\nabla f(\mathbf{x}_t)\|_2^2$$
 not true for constrained case

• Step 2: by convexity,

$$f(\mathbf{x}_{t+1}) \leq f(\mathbf{x}^*) + \langle \nabla f(\mathbf{x}_t), \mathbf{x}^* - \mathbf{x}_t \rangle - \frac{1}{2L} \| \nabla f(\mathbf{x}_t) \|_2^2$$

$$= f(\mathbf{x}^*) + \frac{L}{2} \left\{ \| \mathbf{x}_t - \mathbf{x}^* \|_2^2 - \left\| \mathbf{x}_t - \mathbf{x}^* - \frac{1}{L} \nabla f(\mathbf{x}_t) \right\|_2^2 \right\}$$

• Step 3: telescoping

$$f(\mathbf{x}_T) - f(\mathbf{x}^*) = \frac{1}{T} \sum_{t=0}^{T-1} (f(\mathbf{x}_{t+1}) - f(\mathbf{x}^*)) \le \frac{L}{2T} \|\mathbf{x}_0 - \mathbf{x}^*\|^2$$

19/26

Lecture 05 OptML November 1st, 2023

# Convergence Analysis

For the constrained case, we aims to replace  $\nabla f(\mathbf{x})$  in the unconstrained case by

$$g_{\mathcal{C}}(\mathbf{x}) = L(\mathbf{x} - \mathcal{P}_{\mathcal{C}}(\mathbf{x} - \frac{1}{L}\nabla f(\mathbf{x})))$$

We have  $g_{\mathcal{C}}(\mathbf{x}_t) = L(\mathbf{x}_t - \mathbf{x}_{t+1})$ .

• Step 1: descent guarantee

$$f(\mathbf{x}_{t+1}) \leq f(\mathbf{x}_t) - \frac{1}{2L} \|g_{\mathcal{C}}(\mathbf{x}_t)\|_2^2$$

• Step 2:

$$f(\mathbf{x}_{t+1}) \leq f(\mathbf{x}^*) + \langle g_{\mathcal{C}}(\mathbf{x}_t), \mathbf{x}^* - \mathbf{x}_t \rangle - \frac{1}{2L} \|g_{\mathcal{C}}(\mathbf{x}_t)\|_2^2$$

Step 3: telescoping

$$f(\mathbf{x}_T) - f(\mathbf{x}^*) = \frac{1}{T} \sum_{t=0}^{T-1} (f(\mathbf{x}_{t+1}) - f(\mathbf{x}^*)) \le \frac{L}{2T} \|\mathbf{x}_0 - \mathbf{x}^*\|^2$$

20 / 26

#### Outline

Review


2 Projected Gradient Descent

3 Frank-Wolfe Algorithm

Consider following problem:

$$\min_{\mathbf{x}} f(\mathbf{x})$$
  
s.t.  $\mathbf{A}\mathbf{x} \leq \mathbf{b}$ 

Computing projection is very expensive!



#### Algorithm 1 Frank-Wolfe (a.k.a. conditional gradient) Algorithm

Lecture 05 OptML November 1st, 2023 22 / 26

#### Algorithm 2 Frank-Wolfe (a.k.a. conditional gradient) Algorithm

```
for t = 1, 2, ... do
    \mathbf{y}_t = \arg\min_{\mathbf{x} \in \mathcal{C}} \langle \nabla f(\mathbf{x}_t), \mathbf{x} \rangle //direction finding
    \mathbf{x}_{t+1} = (1 - \eta_t)\mathbf{x}_t + \eta_t\mathbf{v}_t
                                                       //line search and update
```

- main step: linearization of the objective function
- appealing when linear optimization is much cheaper than projection
- stepsize:  $\eta_t = \frac{2}{t+2}$

Lecture 05 OptML November 1st. 2023 23 / 26

Let f be convex and *L*-smooth. If  $\eta_t = \frac{2}{t+2}$ , one has

$$f(\mathbf{x}_t) - f(\mathbf{x}^*) \le \frac{2LD^2}{t+2}$$

where  $D = \sup_{\mathbf{x}, \mathbf{y} \in \mathcal{C}} \|\mathbf{x} - \mathbf{y}\|_2$ 

For compact constraint sets, Frank-Wolfe attains  $\varepsilon$ -accuracy with  $O(\frac{1}{\varepsilon})$  iterations.

# Summary

• Frank-Wolfe: projection-free

	stepsize	convergence	iteration
	rule	rate	complexity
convex & smooth problems	$\eta_t symp rac{1}{t}$	$O\left(\frac{1}{t}\right)$	$O\left(\frac{1}{\varepsilon}\right)$

• projected gradient descent

	stepsize rule	convergence rate	iteration complexity
convex & smooth problems	$\eta_t = \frac{1}{L}$	$O\left(\frac{1}{t}\right)$	$O\left(\frac{1}{\varepsilon}\right)$
strongly convex & smooth problems	$\eta_t = \frac{1}{L}$	$O\left(\left(1-\frac{1}{\kappa}\right)^t\right)$	$O\left(\kappa\log\frac{1}{\varepsilon}\right)$

# Questions

